



Berkshire Diabetic Eye Screening Programme

INSTRUCTIONS FOR PATIENTS RECEIVING EYE DROPS DURING DIABETIC EYE SCREENING EXAMINATION

It is quite normal to experience some stinging immediately after the drops are placed in your eyes. Your vision may also be blurred and light may appear brighter than usual for several hours.

(You should **NOT** drive for at least 4 hours after your screening test).

The type of drops you have been given are:

Single use Tropicamide 1% (0.5% for patients who are pregnant) *please delete as appropriate*

Date & time drops given Batch No..... Expiry Date.....

An adverse reaction to these drops is extremely rare. However, if you experience any unusual symptoms such as headache, vomiting, eye pain or continued blurred vision after attending the retinal screening clinic, you should go to your local hospital's eye casualty department or seek advice by calling the following number:

Call: NHS Direct on 111 (24 hours)



Berkshire Diabetic Eye Screening Programme

INSTRUCTIONS FOR PATIENTS RECEIVING EYE DROPS DURING DIABETIC EYE SCREENING EXAMINATION

It is quite normal to experience some stinging immediately after the drops are placed in your eyes. Your vision may also be blurred and light may appear brighter than usual for several hours.

(You should **NOT** drive for at least 4 hours after your screening test).

The type of drops you have been given are:

Single use Tropicamide 1% (0.5% for patients who are pregnant) *please delete as appropriate*

Date & time drops given Batch No..... Expiry Date.....

An adverse reaction to these drops is extremely rare. However, if you experience any unusual symptoms such as headache, vomiting, eye pain or continued blurred vision after attending the retinal screening clinic, you should go to your local hospital's eye casualty department or seek advice by calling the following number:

Call: NHS Direct on 111 (24 hours)